

# Phys 8911 Lecture 6 (M 07.02.2005)

$N=1$  massive irreps

| $s \backslash Y$ | 0 | $\frac{1}{2}$ | 1 | $\frac{3}{2}$ |
|------------------|---|---------------|---|---------------|
| 0                | 2 | 1             |   |               |
| $\frac{1}{2}$    | 1 | 2             | 1 |               |
| 1                |   | 1             | 2 | 1             |
| $\frac{3}{2}$    |   |               | 1 | 2             |
| 2                |   |               |   | 1             |

$N=1$  massless irreps

| $K \backslash h$ | -2 | $-\frac{3}{2}$ | -1 | $-\frac{1}{2}$ | 0 | $\frac{1}{2}$ | 1 | $\frac{3}{2}$ |
|------------------|----|----------------|----|----------------|---|---------------|---|---------------|
| 2                | 1  |                |    |                |   |               |   |               |
| $\frac{3}{2}$    | 1  | 1              |    |                |   |               |   |               |
| 1                |    | 1              | 1  |                |   |               |   |               |
| $\frac{1}{2}$    |    |                | 1  | 1              |   |               |   |               |
| 0                |    |                |    | 1              | 1 |               |   |               |
| $-\frac{1}{2}$   |    |                |    |                | 1 | 1             |   |               |
| -1               |    | 1              | 1  |                |   |               |   |               |
| $-\frac{3}{2}$   | 1  | 1              |    |                |   |               |   |               |
| -2               | 1  |                |    |                |   |               |   |               |
|                  |    |                |    |                | 4 | 4             | 4 | 4             |

"chiral" "vector" multiplets  
 Add CPT conjugates  
 total # states  $(4=d+2)$

$|S\rangle = |m, k\rangle$   
 $\uparrow$  superhelicity  
 $k = 0, \pm \frac{1}{2}, \pm 1, \pm \dots$   
 obtain  $d$  massless particles of helicity  
 $k, k + \frac{1}{2}$   
 $A, -A \rightarrow \text{spin } |A|$

## Comments

1. Massless irreps are "half as big" as the massive irreps

$$\hookrightarrow a_1^+, a_2^+ \rightarrow 4$$

$$\{Q_\alpha, \bar{Q}_{\dot{\alpha}}\} = 2m \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = 2m \delta_{\alpha\dot{\alpha}}$$

$$\{Q_\alpha, Q_\beta\} = 4E \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

2.  $\kappa = 0$  and  $\kappa = \frac{1}{2}$  massless irreps  $\Rightarrow$  same content as  $\kappa = \frac{1}{2}$  massive irrep.

## Extended supersymmetry

Can have more than one spinor generator  $Q_\alpha^A$  where  $A = 1, \dots, N$

$$\text{Obtain } [P^m, Q_\alpha^A] = 0 = [P^m, \bar{Q}_{\dot{\alpha}}^A]$$

$$[M_{ab}, Q_\alpha^A] = (\delta_{ab})_\alpha^\beta Q_\beta^A$$

$$\{Q_\alpha^A, Q_\beta^B\} = 2\delta_{\alpha\beta}^m P_m \delta^A_B$$

However,  $\{Q_\alpha^A, Q_\beta^B\} = \epsilon_{\alpha\beta} X^{AB}$ , where  $X^{AB}$  is internal symmetry, where  $X^{AB} = -X^{BA}$

$$\begin{aligned} \text{Can show } [X^{AB}, Q_\alpha^C] &= [X^{AB}, \bar{Q}_{\dot{\alpha}C}] = [X^{AB}, B^{\dot{\alpha}C}] = \\ &= [X^{AB}, X^{CD}] = 0 \end{aligned}$$

i.e.  $X^{AB}$  commutes with everything  $\rightarrow$   
known as central charges.

see W & B Chap 1.

Note When  $X^{AB} = 0$  the 4d SUSY algebra  
has a  $U(N)$  sym

$$Q_\alpha^A \rightarrow U^A_B Q_\alpha^B, \quad \bar{Q}_{\dot{\alpha}A} \rightarrow \bar{Q}_{\dot{\alpha}B} U^{+B}_A$$

where  $U^A_B$  is a unitary matrix  $N \times N$ .

When  $X^{AB} \neq 0$  this sym is reduced

e.g.  $N=1$ ,  $X^{AB} = 0$  and the  $U(1)$  symmetry  
is the R-symmetry.

### $N > 1$ representations of SUSY

1. Massless states with no central charges

There are  $N$  creation operators

$$a^+_A \quad (A=1, \dots, N)$$

satisfying  $Q, Q$

$$\{a_A, a^+_B\} = \delta_{AB}$$

$$\{a_A, a_B\} = 0 = \{a^+_A, a^+_B\}$$

Start with the ground state  $|0\rangle$  with  
superhelicity  $K$  can form

$$\frac{1}{\sqrt{n!}} a^+_1 \dots a^+_n |0\rangle \quad 1 \leq n \leq N$$

This gives



|                        |     |                    |                |         |                    |
|------------------------|-----|--------------------|----------------|---------|--------------------|
| helicity ( $\lambda$ ) | $k$ | $k + \frac{1}{2}$  | $k + 1$        | $\dots$ | $k + \frac{N}{2}$  |
| # of states            | 1   | $N = \binom{N}{1}$ | $\binom{N}{2}$ | $\dots$ | $\binom{N}{N} = 1$ |

Total # of States  
 $= \sum_{k=0}^N \binom{N}{k} = 2^N$

Note This is not a CPT eigenstate unless  
 $-k = k + \frac{N}{2} \Rightarrow \boxed{k = -\frac{N}{4}}$

So, in general need to include  
 $-k, -k - \frac{1}{2}, \dots, -k - \frac{N}{2}$  helicity states

Examples:  $N=2$

|       |           |   |               |   |           |   |                |    |
|-------|-----------|---|---------------|---|-----------|---|----------------|----|
| $k=0$ | $\lambda$ | 0 | $\frac{1}{2}$ | 1 | CPT conj. | 0 | $-\frac{1}{2}$ | -1 |
|       | # states  | 1 | 2             | 1 |           | 1 | 2              | 1  |

(curved arrows labeled "combine" connect the two sides of the table)

|           |    |                |   |               |   |
|-----------|----|----------------|---|---------------|---|
| $\lambda$ | -1 | $-\frac{1}{2}$ | 0 | $\frac{1}{2}$ | 1 |
| # states  | 1  | 2              | 2 | 2             | 1 |

This is  $N=2$  Vector multiplet.

|                    |           |                |   |               |
|--------------------|-----------|----------------|---|---------------|
| $k = -\frac{1}{2}$ | $\lambda$ | $-\frac{1}{2}$ | 0 | $\frac{1}{2}$ |
|                    | # states  | 1              | 2 | 1             |

$N=2$  hypermultiplet automatically CPT complete Fayet.

$N=4 \quad k=-1$

|           |      |                |     |               |     |
|-----------|------|----------------|-----|---------------|-----|
| $\lambda$ | $-1$ | $-\frac{1}{2}$ | $0$ | $\frac{1}{2}$ | $1$ |
| #states   | 1    | 4              | 6   | 4             | 1   |

$N=4$  vector multiplet  
auto CPT complete

Note 1. Require renormalizability excludes  $|\lambda| > 1$   
 $\Rightarrow N=4$  max possible <sup>(global)</sup> supersymmetry.

~~$N=4$  vector multiplet  
auto CPT complete~~

2. If exclude  $|\lambda| > 2$  then max. amount of SUSY  $N=2$  (local) supersymmetry  $\rightarrow$  supergravity.

2. Massive states - no central charges

There are now  $2N$  creation ops  $(a_a^A)^\dagger$   
 $A=1 \dots N$

satisfying  $\{a_a^A, a_b^B\} = \delta_{ab} \delta_{AB} 1, 2$

There are  $2^{2N} (2Y+1)$  states in each massive irrep.

For each " $N$ " superspin  $Y_{(N)}$  obtain " $N-1$ " superspin

$\left\{ \begin{array}{l} Y_{(N-1)} = \frac{1}{2}, Y_{(N)} = \frac{Y+1}{2} \\ 0, 0, \frac{1}{2} \end{array} \right. \quad Y \neq 0$   
 $\left\{ \begin{array}{l} 0, 0, \frac{1}{2} \\ Y=0 \end{array} \right.$  where  $Y_{(0)} = s = \text{spin of particle}$

Example:  $N=2 \quad Y_{(2)}=0$

